



## Tank Mixing on Kiwifruit >> Preflowering Nordox™ 75WG

The following combinations of products have been tested for compatibility in the spray tank with Nordox 75WG and for plant phytotoxicity in potted plant trials.

The following combinations were shown in the trial to be plant safe on Hayward and Gold 3 leaves.

Therefore we have some confidence that when applied in the field by a well calibrated sprayer, under good drying conditions and without other additives in the spray tank that the results will be similar to our trial work.

The following needs to be noted and adhered to in regard to Nordox 75WG application on Kiwifruit:

- With the specific exception of the products listed in this guide, do not apply with surfactants, foliar fertilisers, oils or other products in the spray tank.
- Do not apply under damp or cold conditions that may result in slow drying.
- Ensure sprayer is well calibrated, coverage is even and water volume does not result in run off.
- Do not apply to stressed or unhealthy vines.

In our trials, Nordox 75WG was applied as per the product label at 37.5g per 100L water. All other product rates as tested, based on product label and a water volume of 1000L per hectare.

These combinations (below) demonstrated both compatibility and crop safety in our trials.

Nordox 75WG + Mesuro1™ (150ml/100L water)

Nordox 75WG + Actigard® (20g/100L water)

Nordox 75WG + Prodigy™ (50mls/100L water)

Nordox 75WG + Comic™ (8.6g/100L water)

Nordox 75WG + Actigard® (20g/100L water)  
+ Comic (8.6g/100L water)

Nordox 75WG + Gro-Wet™ (125ml/100L water)\*

Nordox 75WG + Bio-Bit™ (50g/100L water)

Nordox 75WG + Boost-It™ (0.5kg/100L water)

Nordox 75WG + Alga 600™ (0.1kg/100L water)

\* Gro-Wet only when used as per the label for concentrate spraying.

Please note the following were either found to be not plant safe in our trial, or known to be unsafe practices.

- Tank mixture of Nordox 75WG and Urea was phytotoxic.
- Tank mixture of Nordox 75WG and Spraying Oil was phytotoxic.
- Application of Spraying Oil within 7 days of a Nordox 75WG application can result in phytotoxicity.
- Tank mixing any acidifying product with Nordox 75WG or any product that may enhance absorption into the leaf (e.g. Amino acid based products) will enhance phytotoxicity.

Prodigy™ is a Trademark of Dow Agrosiences.  
Bio-Bit™ is a trademarks of Valent Biosciences.  
Mesuro1™ is a trademarks of Bayer Crop Sciences.  
Actigard® is a registered trademarks of a Syngenta Group Company.

Disclaimer: Canopy damage is possible from other sprays, plant health and due to weather conditions such as hail or wind. Formulations of chemistry from other suppliers may change without the knowledge of AgriNova NZ Ltd and may result in differing responses. This document is given as a guide and combinations should be checked on your orchard if uncertainty exists. As no control can be exercised over the manner in which or the conditions under which this product is used, all conditions and warranties expressed or implied, other than those required by statute are excluded. No responsibility will be accepted for any damage or injury whatsoever arising from the storage, handling, application or use of this product.

NORDOX 75WG

